arXiv:1209.1939v2 [physics.atom-ph] 18 Oct 2012

Collapse-revival dynamics in strongly laser-driven electrons

O. D. Skoromnik,[1] I. D. Feranchuk,[2] and C. H. Keitel[1]
[1]*Max Planck Institute for Nuclear Physics, Saupfercheckweg 1, 69117 Heidelberg, Germany*
[2]*Belarusian State University, 4 Nezavisimosty Ave., 220030, Minsk, Belarus*

The relativistic quantum dynamics of an electron in an intense single-mode quantized electromagnetic field is investigated with special emphasis on the spin degree of freedom. In addition to fast spin oscillations at the laser frequency, a second time scale is identified due to the intensity dependent emissions and absorptions of field quanta. In analogy to the well-known phenomenon in atoms at moderate laser intensity, we put forward the conditions of collapses and revivals for the spin evolution in laser-driven electrons starting at feasible 10^{18} W/cm^2.

PACS number(s): 12.20.-m, 13.88.+e

The motion of a relativistic electron in the field of a plane electromagnetic wave was first evaluated by Volkov [1]. While the solution of this problem initiated numerous investigations of quantum electrodynamical (QED) effects in the presence of a strong electromagnetic field, the fundamental derivations of this part of QED were developed in a series of papers [2–4] about fifty years ago. Nevertheless, with the development of powerful lasers providing intensities up to $I \sim 10^{22}$W/cm^2 [5] and ambitions plans beyond [6], the analysis of various quantum phenomena [7] such as multi-photon Compton scattering [8–10], electron–positron pair creation [11], spin effects [12], and quantum plasmas [13] has recently become increasingly relevant. In all this research, however, the external electromagnetic field is considered as classical, and its quantum fluctuations are generally believed to be negligible for interactions with very strong fields.

At the same time, it is well known from quantum optics that the dynamics of an atom in a comparably weaker and resonant laser field depends on the quantum fluctuations of the field. One of the most interesting phenomena of this kind is the collapse–revival effect in the evolution of the Jaynes–Cummings model [14, 15] for a two-level atom. This effect was predicted theoretically [16] and later observed experimentally [17]. Its qualitative explanation and analytical description was also given by [18–20]. It was shown that the evolution of the population of the atomic states is characterized by two time scales. The first time scale is the period of the Rabi oscillations while the second slower one is defined by the collapse and revival times of the populations being associated with the absorption and emission of the field quanta.

An electron can on one hand be considered as a two-level system with regard to spin space. On the other hand the electron has no other internal quantum degrees of freedom such as the atom and especially strong laser-electron dynamics is generally considered to be perfectly described within a classical picture. For that reason, the questions of the feasibility of collapse–revival dynamics for laser-driven electrons arises. In this Letter, we present the electron spin evolution in a single-mode quantized field within the framework of Dirac theory. We show that an electron may exhibit collapse and revival dynamics similar to that of two-level atoms and demonstrate a possible implementation of this pure quantum effect already at widely available laser intensities in the moderately relativistic regime.

For an electron in an external electromagnetic field, two different time scales are characteristic [21]. One of these is defined by the frequency ω of the electromagnetic field, i.e. in natural units $\hbar = c = 1$ as used in the complete article: $T_0 = 2\pi/\omega \sim 1.2 \cdot 10^{-4}$cm, for the typical photon energy $\omega = 1$eV. The other scale is called coherence time T_c [2] and is defined by the time required to induce a 2π phase shift in the Volkov state of the electron.

The existence of these two time scales is expected to be simultaneously observable when the total interaction time of the electron in the laser field T_{i} satisfies

$$T_{\mathrm{i}} > T_c \gg T_0,$$

which, as shown later, may indeed be fulfilled for realistic parameters of the laser pulse.

In a more general view, the slow oscillations are characterized by the radiation shift of the electron's quasi-energy, which is defined via the Dirac equation with mass operator [22]. The shift appears both due to the quantum nature of the driving electromagnetic field and the environmental vacuum field. However, the laser-driven processes of emission and absorption of photons with frequency ω are the main contribution to the transition amplitudes that are proportional to the number of quanta in the radiation mode [23]. Therefore, the system of an electron and a one mode quantized field can be investigated as a closed system in a non-perturbative way.

We benefit from the fact that the state vector of a system with definite total quasi-momentum in a single-mode quantized field can be found exactly. For the first time, the exact solution of the Dirac equation

$$\left(i\gamma^{\mu}\partial_{\mu} - \gamma^{\mu} e A_{\mu} - m\right)\psi = 0, \tag{1}$$

with the potential containing the only significant mode [23]

$$A_{\mu} = \frac{e_{\mu}}{\sqrt{2\omega V}}\left(a e^{-i(kx)} + a^{\dagger} e^{i(kx)}\right), \tag{2}$$

Dirac matrices γ^{μ}, polarization vector e^{μ}, normalization volume V, photon four-vector k, photon annihilation and creation operators a and $a^\dagger$ of the laser mode, electron charge e and mass m, was found by Berson [24], via the Bargmann representation of the creation and annihilation operators. This solution, however, is not convenient for our calculation, and we obtained correspondingly the exact solution of Equation (1) in operator form, reading

$$\psi_{qn} = \frac{1}{\sqrt{2\varepsilon_n}} e^{-i(qx)+ia^\dagger a(kx)} \left(1 + \frac{\hat{k}\hat{b}}{2(q\cdot k)}(a+a^\dagger)\right) \cdot e^{\alpha(a^\dagger - a)} e^{-\frac{\eta}{2}(a^2 - a^{\dagger 2})} u(p_n)|n\rangle, \tag{3}$$

with $\hat{f} = \gamma^\mu f_\mu$ for any four-vector f, the total four-momentum of the system of the electron and the field q, $b^\mu = ee^\mu/\sqrt{2V\omega}$, $\alpha = -(q\cdot b)/((q\cdot k) - 2b^2)$, $\mathrm{ch}\eta = 1/2(\sqrt{\varkappa} + 1/\sqrt{\varkappa})$, $\varkappa = 1/\sqrt{1 - 2b^2/(q\cdot k)}$, the quasi-momentum of the electron p_n, and $\varepsilon_n = \sqrt{p_n^2 + m^2}$. The quasi-momentum of the electron depends on the field quantum number, yielding

$$p_n = q - k\left(\sqrt{1 - \frac{2b^2}{(q\cdot k)}}(n + \frac{1}{2}) - \frac{1}{2} - - \frac{(q\cdot b)^2}{(q\cdot k)} \frac{1}{(q\cdot k) - 2b^2}\right). \tag{4}$$

The set of solutions (3) is orthogonal and complete, i. e.

$$\frac{1}{(2\pi)^3} \int \psi^\dagger_{q'n'} \psi_{qn} d\boldsymbol{r} = 2\varepsilon_n \delta_{n'n} \delta(\boldsymbol{q}' - \boldsymbol{q}). \tag{5}$$

In order to demonstrate that an electron in a single-mode quantized field evolves in a similar way as an atomic two-level system, we consider the observable value of the electron spin:

$$s^\mu(\boldsymbol{x}, t) = \frac{\langle\psi|\gamma^0\gamma^5\gamma^\mu\delta(\boldsymbol{x} - \boldsymbol{r}')|\psi\rangle}{\langle\psi|\psi\rangle}. \tag{6}$$

Before the electron crosses the boundary of the light beam at $t = 0$, it is described by the free solution of the Dirac equation

$$\psi_0 = e^{i\boldsymbol{p}_0(\boldsymbol{r} - \boldsymbol{r}_0)} \frac{u(p_0)}{\sqrt{2\varepsilon_0}}, \tag{7}$$

and the field is in the coherent state

$$|\beta\rangle = \sum_{n=0}^{\infty} \frac{\beta^n}{\sqrt{n!}} |n\rangle e^{-\beta^2/2}, \tag{8}$$

with the average number of photons $\bar{n} = \beta^2$.

The variable $\boldsymbol{r}_0$ defines the position of the electron where we have to take into account that in a real experiment many electrons are distributed in a bunch. As a consequence we will later average over this variable by integrating over the appropriate density [25, 26].

During the interaction the system of the electron and the field is described by the wave function which is a linear combination of the exact solutions (3), i. e.

$$\Psi(x) = \int d\boldsymbol{q} \sum_n C_{\boldsymbol{q},n} \psi_{qn}(x); \; x = (t, \boldsymbol{x}). \tag{9}$$



Figure 1. (Color online) Electron motion in a single-mode quantized field without momentum change (coherent scattering). $\boldsymbol{p}_0$ is the initial momentum of the electron, $\boldsymbol{b}$ the polarization vector of the field, the z axis is directed along the wave vector $\boldsymbol{k}$, $\boldsymbol{z}_\perp$ denotes the plane perpendicular to z, a and $a^\dagger$ are the annihilation and creation operators, respectively, ω is the frequency of field, V a normalization volume, and e is the electron charge.

After crossing the border of the field, the interaction holds for $t_0 \sim \frac{d}{v_0 \sin\theta}$, where d is the "thickness" of the laser pulse (see Fig. 1), $\boldsymbol{v}_0$ the injection velocity of the electron, and θ is the angle between $\boldsymbol{k}$ and $\boldsymbol{v}_0$.

At $t = t_0$, the interaction is turned off, the electron becomes free, and the detector may measure the spin $\boldsymbol{s}(t_0)$. To guarantee that only the spin degree of freedom changes, we will analyse the evolution for the polarization state of electrons only with the same momentum as in the primary beam. This case is essentially different from the problem of the polarization of an electron as a result of Compton scattering [8].

A change in the interaction time t_0 will lead to different spin values. This time can be changed in two ways. The first is a change in the initial energy of the electron. The larger the electron energy is, the less time it spends in the field. The second way is a change in the interaction time, changing the angle θ. The closer the angle θ is to 0 or π, the more time the electron interacts with the field.

Here, we investigate the case of a strong field such that the average number of photons $\bar{n} = \beta^2$ is large. Therefore, the main contributions to the spin value (6) will arise from $n \gg 1$ and the coefficients $C_{\boldsymbol{q},n}$ can be evaluated in this limit. This allows us to calculate the average value of the spin analytically. Let us consider here the outline of the calculation only.

From the initial condition

$$e^{i\boldsymbol{p_0}(\boldsymbol{r} - \boldsymbol{r}_0)} \frac{u(p_0)}{\sqrt{2\varepsilon_0}} |\beta\rangle = \Psi\big|_{t=0}, \tag{10}$$

the unknown coefficients $C_{\boldsymbol{q},n}$ in the limit of large n were found. Then the wave function (9) was inserted into the spin definition (6) and the matrix elements between the field states $T_{n',n}$ were calculated. Actually, these matrix elements were evaluated using the cumulant method [27], which corresponds to a non-perturbative partial summation of the infinite series. The integrals over $\boldsymbol{q}$, $\boldsymbol{q}'$, and the summation over the field quantum number n were carried out analytically. The evaluation of the summation over the field quantum number n was performed analogously to [18] by means of the saddle point method. The saddle point n_0 has quadratic and linear terms in β. The zero order approximation $n_0 = \beta^2$ gives the quasi-classical Volkov limit, and the first order corrections which are proportional to $\beta\alpha$ induce the quantum fluctuations of interest here. In the last step, an averaging over the initial electron coordinate $\boldsymbol{r}_0$ was carried out and the matrix element in spinor space was calculated. As a result, the dependence of the spin on time was found in the form

$$\begin{aligned} \langle s^\mu(t)\rangle &= \frac{m}{\varepsilon_0}a_0^\mu - \frac{m}{\varepsilon_0}k^\mu(a_0\cdot k)\frac{\beta^2b^2}{(p_0\cdot k)^2}\left(1+\mathrm{Re}\,\Pi_2\right) \\ &+\left[\frac{m}{\varepsilon_0}\frac{\beta}{(p_0\cdot k)}\left(k^\mu(a_0\cdot b) - b^\mu(a_0\cdot k)\right)\right]2\mathrm{Re}\,\Pi_1, \\ \Pi_l &= (-i)^l e^{i\omega l t\left(1-\frac{p_{0z}}{\varepsilon_0}\right)} J_l(4\alpha\beta\sin\frac{\omega^2 l t}{2\varepsilon_0}), \quad l\in\{1,2\}, \end{aligned} \tag{11}$$

where J_l is the Bessel function of order l and a_0 an initial spin four-vector of the electron. This value of the spin may be compared with the time averaged Volkov one, i. e.

$$\langle\bar{\boldsymbol{s}}\rangle = \boldsymbol{a}_0\frac{m}{\varepsilon} - \frac{me^2}{2\varepsilon(k\cdot p)^2}\boldsymbol{k}(k\cdot a_0)\bar{A^2}, \tag{12}$$

where the bar on top of a symbol denotes the averaging in time and A represents the four-potential of the classical field. Actually, this averaging corresponds to that over the initial phases of the electron in the beam when entering the space with the field.

For the analysis of the evolution of the electron polarization, we also need to calculate the electron current density $j^\mu(\boldsymbol{x},t) = \langle\psi|\gamma^\mu\delta(\boldsymbol{x}-\boldsymbol{r}')|\psi\rangle/\langle\psi|\psi\rangle$ averaged over the electron initial coordinate $\boldsymbol{r}_0$. This value can be evaluated in the same way as considered above for the electron spin density, yielding

$$\begin{aligned} j^\mu(t) &= \frac{p^\mu}{\varepsilon_0} - \frac{\beta b^\mu}{\varepsilon_0}2\mathrm{Re}\Pi_1 \\ &+\frac{k^\mu}{\varepsilon_0}\left(\frac{\beta(b\cdot p)}{(k\cdot p)}2\mathrm{Re}\Pi_1 - \frac{\beta^2b^2}{(k\cdot p)}\left(1+\mathrm{Re}\,\Pi_2\right)\right). \end{aligned} \tag{13}$$

The electron polarization is characterized by its helicity—the projection of the spin $\boldsymbol{s}$ onto the direction of its momentum $\boldsymbol{\nu} = \boldsymbol{p}_0/p_0$. The eigenvalues of the helicity operator for the free electron are ±1. In accordance with the general rules of quantum mechanics, the expectation value of the helicity operator can be written as

$$\langle\boldsymbol{\Sigma}\cdot\boldsymbol{\nu}\rangle = p_1\cdot 1 + p_{-1}\cdot(-1), \tag{14}$$

with p_1 and p_{-1} being the probabilities of observing the electron with helicity $+1$ and -1, respectively. Usually, one considers at the beginning the situation of p_1 being unity and p_{-1} zero, and the interest is how p_1 and p_{-1} are modified due to the interaction.

However, in real experiments, not a single electron, but rather a beam of electrons is considered with N being the number of electrons in the beam. If the initial electron beam was fully polarized, some quantity of electrons with opposite polarization should appear after the interaction is finished (see Fig. 1). Namely, the number N_{-1} of such electrons would be equal to $p_{-1}N$ if in the initial moment of time $t = 0$ the helicity of the electron was $+1$.

To calculate the final polarization, we choose the coordinate system with z-axis directed along $\boldsymbol{k}$, an initial electron momentum $\boldsymbol{p}_0$ displaying an angle θ with the z-axis, and a field being linearly polarized and directed perpendicular to the z-axis (Fig. 1). We assume that at the initial moment of time the vector $\boldsymbol{\zeta}$ being the average spin in the electron rest frame is directed along the momentum in the laboratory frame. As a consequence $\boldsymbol{p}\cdot\boldsymbol{\zeta} = p$, which corresponds to a helicity of the electron of $+1$.

In order to find the polarization of the electron after the interaction with the field, we project the spin vector $\boldsymbol{s}$ on the electron momentum direction $\boldsymbol{\nu} = (0, \sin\theta, \cos\theta)$ and normalize it via the probability of finding an electron at the observation point, i.e., dividing by $j^0(t)$, the zeroth component of the current density vector, defined in (13). Using this fact, Equation (14) and the condition that $p_1 + p_{-1} = 1$, the probabilities to find the electron in the transmitted and thus non scattered beam in various polarization states can be calculated as

$$\begin{aligned} p_1 &= \frac{1}{2} + \frac{1}{2}\frac{1+\xi^2 f - 2\xi g}{1+\xi^2 f_1 - 2\xi g}, \\ p_{-1} &= \frac{1}{2} - \frac{1}{2}\frac{1+\xi^2 f - 2\xi g}{1+\xi^2 f_1 - 2\xi g}, \end{aligned} \tag{15}$$

where $f = (v_0 - \cos\theta)\cos\theta/(1-v_0\cos\theta)^2\left(1+\mathrm{Re}\ \Pi_2\right)$, $g = v_0\sin\theta/(1-v_0\cos\theta)\mathrm{Re}\ \Pi_1$, $f_1 = \left(1+\mathrm{Re}\ \Pi_2\right)/(1-v_0\cos\theta)$, and the dimensionless parameter $\xi = \beta b/\varepsilon_0$ was introduced.

These probabilities may be compared with the Volkov probabilities written in the same variables, i. e.

$$p_1 = \frac{1}{2} + \frac{1}{2}\frac{1+\xi^2 f_v}{1+\xi^2 f_{1v}}, \quad p_{-1} = \frac{1}{2} - \frac{1}{2}\frac{1+\xi^2 f_v}{1+\xi^2 f_{1v}}, \tag{16}$$

where $f_v = (v_0 - \cos\theta)\cos\theta/(1-v_0\cos\theta)^2$ and $f_{1v} = 1/(1-v_0\cos\theta)$.

One can see that contrary to the Volkov case, the fully quantum probabilities depend on time. This means that the quantum fluctuations can change the system dynamics.

Let us find the parameter which governs the slow oscillations in the time evolution in (15). For this purpose, we investigate the system's evolution for small times t. In this case, the sine inside the Bessel functions can be expanded in its Taylor series and one can find that the



Figure 2. (Color online) The probability of finding the electron with flipped polarization as a function of the angle θ for an intensity $I = 10^{18}\mathrm{W/cm}^2$, a pulse duration $\tau = 1.5 \cdot 10^{-3}\mathrm{cm}$ corresponding to 50 fs, a frequency of the photon $\omega = 7.8 \cdot 10^4\mathrm{cm}^{-1}$, an initial probability $p_{-1} = 0$, and γ values of the electron equal to 5 and 10.

amplitude of the quantum fluctuations is defined via the parameter ξ which corresponds to x in [4].

Modern lasers can reach nowadays high intensities [5] up to $10^{22}\mathrm{W/cm}^2$ with a pulse duration of about 30 fs. For our concrete analysis, we choose as intensity $I = 10^{18}\mathrm{W/cm}^2$, as pulse duration $\tau = 1.5 \cdot 10^{-3}\mathrm{cm}^{-1}$ corresponding to 50 fs, and as photon frequency $\omega = 7.8 \cdot 10^4\mathrm{cm}^{-1}$, i. e. a wavelength of 800 nm.

As was already mentioned, the interaction time can be adapted either via the electron's energy or by changing the electron's path in the light pulse such as e.g. by varying the entrance angle θ. Fig. 2 displays the probability of finding the particles with the flipped polarization in the electron beam as a function of the equal entrance and exit angles θ between $\boldsymbol{p}_0$ and $\boldsymbol{k}$. As can be seen from the graphs, when the interaction time increases, corresponding to larger angles, characteristic structures of the probabilities appear. In addition to fast oscillations at the frequency of field ω, there are slow oscillations governed by the parameter ξ. These oscillations occur around the mean value, which corresponds to the quasi-classical Volkov case. It should be noted that this special structure appears only when the field is considered as a quantum object: it can not appear in the quasi-classical case.

Fig. 3 shows the dependence of the probability of finding an electron with flipped polarization as a function of the electron energy for the two values of fly-in angle θ equal to 30 and 140 degrees. As in the previous case, there are oscillations which appear due to the quantum nature of the electromagnetic field. The average of these oscillations represents the quasi-classical Volkov probability.

In real experiments an electron beam involves a spread in both initial energy and direction. We ensured that an uncertainty of one percent in energy and in the angular distribution does not change the displayed probabilities in above figures visibly. We emphasise from the order of magnitude in the probabilities in Fig. 2 & 3 that the number of electrons should be well above 1000 and that mutual interactions shall be avoided with appropriate densities.



Figure 3. (Color online) The probability of finding an electron with flipped polarization as a function of γ for an intensity $I = 10^{18}\mathrm{W/cm}^2$, a pulse duration $\tau = 1.5 \cdot 10^{-3}\mathrm{cm}$ corresponding to 50 fs, a frequency of the photon $\omega = 7.8 \cdot 10^4\mathrm{cm}^{-1}$, an initial probability $p_{-1} = 0$, and values of the fly-in angle θ equal to 30 and 140 degrees.

Note that the atomic two-level system possesses an analogous behaviour. In this case, the level population is considered instead of the electron polarization. The fast oscillations in the population inversion essentially depend on the Rabi frequency while for an electron in a quantized field, it can be linked to the oscillations at the frequency of the electromagnetic field ω. The structure of the slow oscillations is reminiscent of the "collapse–revival" effect for inverted populations, according to which the population inversion of the two-level system vanishes but after some time revives again. This effect is purely quantum mechanical and can not be found in a quasi-classical analysis. We stress in concluding, that the presented quantum results coincide with those via the quasi-classical Volkov solution if the parameter $\beta\alpha$ is set equal to zero.

The authors are grateful to Dr. K. Z. Hatsagortsyan, Dr. A. Di Piazza and S. Meuren for useful discussions.

---

[1] D. M. Volkov, JETP **7**, 1286 (1937).
[2] A. I. Nikishov and V. I. Ritus, Sov. Phys. JETP **19**, 529 (1964); I. I. Goldman, *ibid.* **19**, 954 (1964); N. B. Narozhny, A. I. Nikishov, and V. I. Ritus, *ibid.* **20**, 622 (1965).
[3] H. R. Reiss, Journal of Mathematical Physics **3**, 59

(1962).

[4] V. I. Ritus, Journal of Russian Laser Research **6**, 497 (1985), 10.1007/BF01120220.

[5] V. Bagnoud, B. Aurand, A. Blazevic, S. Borneis, C. Bruske, B. Ecker, U. Eisenbarth, J. Fils, A. Frank, E. Gaul, S. Goette, C. Haefner, T. Hahn, K. Harres, H.-M. Heuck, D. Hochhaus, D. Hoffmann, D. Javorková, H.-J. Kluge, T. Kuehl, S. Kunzer, M. Kreutz, T. Merz-Mantwill, P. Neumayer, E. Onkels, D. Reemts, O. Rosmej, M. Roth, T. Stoehlker, A. Tauschwitz, B. Zielbauer, D. Zimmer, and K. Witte, Applied Physics B: Lasers and Optics **100**, 137 (2010); E. Gerstner, Nature **446**, 16 (2007); V. Yanovsky, V. Chvykov, G. Kalinchenko, P. Rousseau, T. Planchon, T. Matsuoka, A. Maksimchuk, J. Nees, G. Cheriaux, G. Mourou, and K. Krushelnick, Opt. Express **16**, 2109 (2008).

[6] ELI, Extreme Light Infrastructure, http://www.extreme-light-infrastructure.eu/ (2011); HiPER, High Power Laser Energy Research, http://www.hiper-laser.org/ (2011).

[7] A. Di Piazza, C. Müller, K. Z. Hatsagortsyan, and C. H. Keitel, Rev. Mod. Phys. **84**, 1177 (2012); F. Ehlotzky, K. Krajewska, and J. Z. Kamiński, Reports on Progress in Physics **72**, 046401 (2009); G. A. Mourou, T. Tajima, and S. V. Bulanov, Rev. Mod. Phys. **78**, 309 (2006).

[8] D. V. Karlovets, Phys. Rev. A **84**, 062116 (2011).

[9] F. Mackenroth and A. Di Piazza, Phys. Rev. A **83**, 032106 (2011); M. Boca and A. Oprea, Physica Scripta **83**, 055404 (2011); D. Seipt and B. Kämpfer, *ibid.* **83**, 022101 (2011); M. Boca and V. Florescu, *ibid.* **80**, 053403 (2009); C. Harvey, T. Heinzl, and A. Ilderton, *ibid.* **79**, 063407 (2009).

[10] M. Boca, V. Florescu, and M. Gavrila, Phys. Rev. A **85**, 023425 (2012); J. Bergou and S. Varro, Journal of Physics A: Mathematical and General **14**, 2281 (1981).

[11] A. I. Titov, H. Takabe, B. Kämpfer, and A. Hosaka, Phys. Rev. Lett. **108**, 240406 (2012); T.-O. Müller and C. Müller, Physics Letters B **696**, 201 (2011); A. Di Piazza, A. I. Milstein, and C. Müller, Phys. Rev. A **82**, 062110 (2010).

[12] S. Meuren and A. Di Piazza, Phys. Rev. Lett. **107**, 260401 (2011); F. H. M. Faisal and S. Bhattacharyya, *ibid.* **93**, 053002 (2004); M. W. Walser, D. J. Urbach, K. Z. Hatsagortsyan, S. X. Hu, and C. H. Keitel, Phys. Rev. A **65**, 043410 (2002).

[13] F. Haas, B. Eliasson, and P. K. Shukla, Phys. Rev. E **85**, 056411 (2012); G. Brodin, A. P. Misra, and M. Marklund, Phys. Rev. Lett. **105**, 105004 (2010).

[14] E. T. Jaynes and F. W. Cummings, Proc. Inst. Elect. Eng. **51** (1963); F. W. Cummings, Phys. Rev. **140**, A1051 (1965).

[15] B. W. Shore and P. L. Knight, Journal of Modern Optics **40**, 1195 (1993).

[16] J. H. Eberly, N. B. Narozhny, and J. J. Sanchez-Mondragon, Phys. Rev. Lett. **44**, 1323 (1980); N. B. Narozhny, J. J. Sanchez-Mondragon, and J. H. Eberly, Phys. Rev. A **23**, 236 (1981); G. Rempe and H. Walther, *ibid.* **42**, 1650 (1990); M. Fleischhauer and W. P. Schleich, *ibid.* **47**, 4258 (1993).

[17] G. Rempe, H. Walther, and N. Klein, Phys. Rev. Lett. **58**, 353 (1987); M. Brune, F. Schmidt-Kaler, A. Maali, J. Dreyer, E. Hagley, J. M. Raimond, and S. Haroche, *ibid.* **76**, 1800 (1996).

[18] I. D. Feranchuk and A.V.Leonov, Physics Letters A **373**, 517 (2009); I. D. Feranchuk and A. Leonov, *ibid.* **375**, 385 (2011); I. D. Feranchuk and A. V. Leonov, *ibid.* **373**, 4113 (2009).

[19] M. O. Scully and M. S. Zubairy, *Quantum Optics* (Cambridge University Press, Cambridge, 1997).

[20] P. Filipowicz, Journal of Physics A: Mathematical and General **19**, 3785 (1986).

[21] Y. B. Zelídovich, Sov. Phys. Usp **18**, 79 (1975).

[22] A. I. Akhiezer and V. B. Beresteckij, *Quantum Electrodynamics*, 3rd ed. (Nauka, 1969).

[23] L. D. Landau and E. M. Lifschitz, *Quantum Electrodynamics*, 1st ed. (Pergamon, 1971), see paragraph 101.

[24] I. Berson, Soviet Physics JETP **29**, 871 (1969); see also P. Filipowicz and J. Mostowski, Physics Letters A **86**, 356 (1981); P. Filipowicz, Journal of Physics A: Mathematical and General **18**, 1675 (1985).

[25] M. L. Goldberger and K. M. Watson, *Collision Theory*, 1st ed. (John Wiley & Sons, 1964).

[26] L. D. Landau and E. M. Lifschitz, *Quantum Mechanics*, 2nd ed. (Pergamon, 1965).

[27] R. Kubo, JPSJ **17**, 1100 (1962).

[28] V. G. Baryshevskii and A. G. Shekhtman, Physics of Atomic Nuclei **57**, 1299 (1994).

[29] R. P. Feynman, *Statistical Mechanics A Set of Lectures* (Addison-Wesley, 1972).

[30] I. S. Gradshtein and I. M. Ryzhik, *Tables of integrals, sums, series and products* (FIZMATGIZ, Moscow, 1963).

[31] L. D. Landau and E. M. Lifschitz, *Statistical physics*, 3rd ed. (Nauka, 1976).

[32] P. M. Morse and H. Feshbach, *Methods of Theoretical Physics* (McGraw-Hill, New York, 1953).